What You Can do if You are at Higher Risk of Severe Illness from COVID-19

(Print Resources Web Page: https://www.cdc.gov/coronavirus/2019-ncov/communication/print-resources.html)

Are You at Higher Risk for Severe Illness?



Based on what we know now, those at high-risk for severe illness from COVID-19 are:

- People aged 65 years and older
- · People who live in a nursing home or long-term care facility

People of all ages with underlying medical conditions, particularly if not well controlled, including:

- People with chronic lung disease or moderate to severe asthma
- · People who have serious heart conditions
- People who are immunocompromised
- Many conditions can cause a person to be immunocompromised, including cancer treatment, smoking, bone marrow or organ transplantation, immune deficiencies, poorly controlled HIV or AIDS, and prolonged use of corticosteroids and other immune weakening medications.
- People with severe obesity (body mass index [BMI] of 40 or higher)
- · People with diabetes
- · People with chronic kidney disease undergoing dialysis
- People with liver disease

Here's What You Can do to Help Protect Yourself



Stay home if possible.



Wash your hands often.



Avoid close contact (6 feet, which is about two arm lengths) with people who are sick.



Clean and disinfect frequently touched surfaces.



Avoid all cruise travel and non-essential air travel.

Call your healthcare professional if you are sick.

For more information on steps you can take to protect yourself, see CDC's How to Protect Yourself.




Symptoms of Coronavirus (COVID-19)

Know the symptoms of COVID-19, which can include the following:















Symptoms can range from mild to severe illness, and appear 2-14 days after you are exposed to the virus that causes COVID-19.

*Seek medical care immediately if someone has emergency warning signs of COVID-19.

- Trouble breathing
- Persistent pain or pressure in the chest
- New confusion

- Inability to wake or stay awake
- Bluish lips or face

This list is not all possible symptoms. Please call your medical provider for any other symptoms that are severe or concerning to you.




How to Safely Wear and Take Off a Cloth Face Covering

Accessible: https://www.cdc.gov/coronavirus/2019-ncov/prevent-getting-sick/diy-cloth-face-coverings.html

WEAR YOUR FACE COVERING CORRECTLY

- · Wash your hands before putting on your face covering
- Put it over your nose and mouth and secure it under your chin
- · Try to fit it snugly against the sides of your face
- · Make sure you can breathe easily
- Do not place a mask on a child younger than 2








USE THE FACE COVERING TO HELP PROTECT OTHERS

- Wear a face covering to help protect others in case you're infected but don't have symptoms
- Keep the covering on your face the entire time you're in public
- Don't put the covering around your neck or up on your forehead
- · Don't touch the face covering, and, if you do, clean your hands

FOLLOW EVERYDAY HEALTH HABITS

- · Stay at least 6 feet away from others
- Avoid contact with people who are sick
- Wash your hands often, with soap and water, for at least 20 seconds each time
- Use hand sanitizer if soap and water are not available





TAKE OFF YOUR CLOTH FACE COVERING CAREFULLY, WHEN YOU'RE HOME

- Untie the strings behind your head or stretch the ear loops
- Handle only by the ear loops or ties
- Fold outside corners together
- · Place covering in the washing machine
- · Wash your hands with soap and water




Cloth face coverings are not surgical masks or N-95 respirators, both of which should be saved for health care workers and other medical first responders.

For instructions on making a cloth face covering, see:

Stop Germs! Wash Your Hands.

When?

- After using the bathroom
- · Before, during, and after preparing food
- Before eating food
- Before and after caring for someone at home who is sick with vomiting or diarrhea
- After changing diapers or cleaning up a child who has used the toilet
- · After blowing your nose, coughing, or sneezing
- After touching an animal, animal feed, or animal waste
- · After handling pet food or pet treats
- After touching garbage



How?



Wet your hands with clean, running water (warm or cold), turn off the tap, and apply soap.



Lather your hands by rubbing them together with the soap. Be sure to lather the backs of your hands, between your fingers, and under your nails.



Scrub your hands for at least 20 seconds. Need a timer? Hum the "Happy Birthday" song from beginning to end twice.



Rinse hands well under clean, running water.



Dry hands using a clean towel or air dry them.

Keeping hands clean is one of the most important things we can do to stop the spread of germs and stay healthy.



www.cdc.gov/handwashing



Prevent the spread of COVID-19 if you are sick

Accessible version: https://www.cdc.gov/coronavirus/2019-ncov/if-you-are-sick/steps-when-sick.html

If you are sick with COVID-19 or think you might have COVID-19, follow the steps below to care for yourself and to help protect other people in your home and community.

Stay home except to get medical care.

 Stay home. Most people with COVID-19 have mild illness and are able to recover at home without medical care. Do not leave your home, except to get medical care. Do not visit public areas.



- Take care of yourself. Get rest and stay hydrated. Take overthe-counter medicines, such as acetaminophen, to help you feel better.
- Stay in touch with your doctor. Call before you get medical care. Be sure to get care if you have trouble breathing, or have any other emergency warning signs, or if you think it is an emergency.
- · Avoid public transportation, ride-sharing, or taxis.

Separate yourself from other people and pets in your home.

- As much as possible, stay in a specific room and away from other people and pets in your home. Also, you should use a separate bathroom, if available. If you need to be around other people or animals in or outside of the home, wear a cloth face covering.
 - See COVID-19 and Animals if you have questions about pets: https://www.cdc.gov/coronavirus/2019-ncov/faq. html#COVID19animals
 - Additional guidance is available for those living in close quarters. (https://www.cdc.gov/coronavirus/2019-hj ncov/ daily-life-coping/living-in-close-quarters.html) and shared housing (https://www.cdc.gov/coronavirus/2019-ncov/ daily-life-coping/shared-housing/index.html).

Monitor your symptoms.

 Symptoms of COVID-19 include fever, cough, and shortness of breath but other symptoms may be present as well.



 Follow care instructions from your healthcare provider and local health department. Your local health authorities will give instructions on checking your symptoms and reporting information.

When to Seek Emergency Medical Attention

Look for **emergency warning signs*** for COVID-19. If someone is showing any of these signs, **seek emergency medical care immediately:**

- Trouble breathing
- Persistent pain or pressure in the chest
- New confusion
- Bluish lips or face
- · Inability to wake or stay awake

*This list is not all possible symptoms. Please call your medical provider for any other symptoms that are severe or concerning to you.

Call 911 or call ahead to your local emergency facility: Notify the operator that you are seeking care for someone who has or may have COVID-19.

Call ahead before visiting your doctor.

- Call ahead. Many medical visits for routine care are being postponed or done by phone or telemedicine.
- If you have a medical appointment that cannot be postponed, call your doctor's office, and tell them you have or may have COVID-19.

If you are sick, wear a cloth covering over your nose and mouth.

 You should wear a cloth face covering over your nose and mouth if you must be around other people or animals, including pets (even at home).



- You don't need to wear the cloth face covering if you are alone.
 If you can't put on a cloth face covering (because of trouble
 breathing for example), cover your coughs and sneezes in some
 other way. Try to stay at least 6 feet away from other people.
 This will help protect the people around you.
- Cloth face coverings should not be placed on young children under age 2 years, anyone who has trouble breathing, or anyone who is not able to remove the covering without help.

Note: During the COVID-19 pandemic, medical grade facemasks are reserved for healthcare workers and some first responders. You may need to make a cloth face covering using a scarf or bandana.



Cover your coughs and sneezes.

- Cover your mouth and nose with a tissue when you cough or sneeze.
- Throw used tissues in a lined trash can.
- Immediately wash your hands with soap and water for at least 20 seconds. If soap and water are not available, clean your hands with an alcohol-based hand sanitizer that contains at least 60% alcohol.



 Wash your hands often with soap and water for at least 20 seconds. This is especially important after blowing your nose, coughing, or sneezing; going to the bathroom; and before eating or preparing food.



- Use hand sanitizer if soap and water are not available. Use an alcohol-based hand sanitizer with at least 60% alcohol, covering all surfaces of your hands and rubbing them together until they feel dry.
- Soap and water are the best option, especially if your hands are visibly dirty.
- Avoid touching your eyes, nose, and mouth with unwashed hands.

Avoid sharing personal household items.

- Do not share dishes, drinking glasses, cups, eating utensils, towels, or bedding with other people in your home.
- Wash these items thoroughly after using them with soap and water or put them in the dishwasher.

Clean all "high-touch" surfaces everyday.

 Clean and disinfect high-touch surfaces in your "sick room" and bathroom. Let someone else clean and disinfect surfaces in common areas, but not your bedroom and bathroom.



 If a caregiver or other person needs to clean and disinfect a sick person's bedroom or bathroom, they should do so on an as-needed basis. The caregiver/other person should wear a cloth face covering and wait as long as possible after the sick person has used the bathroom.

High-touch surfaces include phones, remote controls, counters, tabletops, doorknobs, bathroom fixtures, toilets, keyboards, tablets, and bedside tables.

- Clean and disinfect areas that may have blood, stool, or body fluids on them.
- Use household cleaners and disinfectants. Clean the area or item with soap and water or another detergent if it is dirty. Then use a household disinfectant.
 - Be sure to follow the instructions on the label to ensure safe and effective use of the product. Many products recommend keeping the surface wet for several minutes to ensure germs are killed. Many also recommend precautions such as wearing gloves and making sure you have good ventilation during use of the product.
 - Most EPA-registered household disinfectants should be effective.

When you can be around others after you had or likely had COVID-19



When you can be around others (end home isolation) depends on different factors for different situations.

I think or know I had COVID-19, and I had symptoms

- You can be with others after
 - 3 days with no fever

AND

symptoms improved

AND

- 10 days since symptoms first appeared
- Depending on your healthcare provider's advice and availability of testing, you might get tested to see if you still have COVID-19. If you will be tested, you can be around others when you have no fever, symptoms have improved, and you receive two negative test results in a row, at least 24 hours apart.

· I tested positive for COVID-19 but had no symptoms

- If you continue to have no symptoms, you can be with others after:
 - 10 days have passed since test
- Depending on your healthcare provider's advice and availability of testing, you might get tested to see if you still have COVID-19. If you will be tested, you can be around others after you receive two negative test results in a row, at least 24 hours apart.
- If you develop symptoms after testing positive, follow the guidance above for "I think or know I had COVID, and I had symptoms."

Handwashing: Keeping Your Family Healthy

Handwashing is an easy, cheap, and effective way to prevent the spread of germs and keep kids and adults healthy. When your family is healthy, you don't have to worry about missing school, work, or other activities.

Help your child develop handwashing skills

Parents and caretakers play an important role in teaching children to wash their hands. Handwashing can become a lifelong healthy habit if you start teaching it at an early age. Teach kids the <u>five easy steps</u> for handwashing—wet, lather, scrub, rinse, and dry—and the key times to wash hands, such as after using the bathroom or before eating. You can find ways to make it fun, like making up your own handwashing song or turning it into a game.

Lead by example

Young children learn by imitating the behaviors of adults in their lives. When you make handwashing part of your routine, you're setting an example for your children to follow.











www.cdc.gov/handwashing



Handwashing can prevent

1in 3 cases of diarrhea



1 in 5

respiratory infections, such as a cold or the flu





Give frequent reminders

Building handwashing skills takes time. At first, your child will need regular reminders of how and when to wash hands. It is especially important to remind children to wash their hands after using the bathroom, before eating, after touching pets, after playing outside, and after coughing, sneezing, or blowing their nose. But once handwashing becomes a habit and a regular part of your child's day, they will practice it throughout their lives.

What if soap and water aren't available?

Washing hands with soap and water is the best way to get rid of germs. If soap and water are not readily available, use an alcohol-based hand sanitizer that has at least 60% alcohol.

Did you know?

Baby wipes may make your hands look clean, but they're not designed to remove germs from your hands. CDC recommends washing hands with soap and water when possible.



Remember to make handwashing a healthy habit at home, school, and at play!